

# Complement Component C4c

**CPT Code:** 86160

**Order Code:** 353

**ABN Requirement:** No

**Specimen:** Serum

**Volume:** 1.0 mL

**Minimum Volume:** 0.5 mL

**Container:** Gel-barrier tube (SST, Tiger Top)

## Collection:

1. Collect and label sample according to standard protocols.
2. Gently invert tube 5 times immediately after draw. DO NOT SHAKE.
3. Allow blood to clot 30 minutes.
4. Centrifuge for 10 minutes.

**Transport:** Store serum at 2°C to 8°C after collection and ship the same day per packaging instructions included with the provided shipping box.

## Stability:

**Ambient (15-25°C):** Not Acceptable

**Refrigerated (2-8°C):** 4 days

**Frozen (-20°C):** 21 days

**Causes for Rejection:** Specimens other than serum; improper labeling; samples not stored properly; samples older than stability limits

**Methodology:** Immunoturbidimetric

**Turn Around Time:** 2 to 3 days

## Reference Range:

Age	Male	Female
<1 Year	Not established	Not established
1-14 Years	14-44 mg/dL	13-46 mg/dL

15-80 Years	15-53 mg/dL	15-57 mg/dL
≥81 Years	Not established	Not established

**Clinical Significance:** Decreased C4 level is associated with acute systemic lupus erythematosus, glomerulonephritis, immune complex disease, cryoglobulinemia, congenital C4 deficiency and generalized autoimmune disease.

*The CPT codes provided are based on AMA guidelines and are for informational purposes only. CPT coding is the sole responsibility of the billing party. Please direct any questions regarding coding to the payer being billed.*